

The structural physical approximations and optimal entanglement witnesses

Kil-Chan Ha¹ and Seung-Hyeok Kye²

¹⁾*Faculty of Mathematics and Statistics,*

Sejong University, Seoul 143-747, Korea

²⁾*Department of Mathematics and Institute of Mathematics,*

Seoul National University, Seoul 151-742, Korea

(Dated: 4 March 2013)

We introduce the notions of positive and copositive types for entanglement witnesses, depending on the distance to the positive part and copositive part. An entanglement witness W is of positive type if and only if its partial transpose W^Γ is of copositive type. We show that if the structural physical approximation of W is separable then W should be of copositive type, and the SPA of W^Γ is never separable unless W is of both positive and copositive type. This shows that the SPA conjecture is meaningful only for those of copositive type. We provide examples to show that the SPA conjecture fails even for the case of copositive types.

PACS numbers: 03.65.Ud, 03.67.Mn

I. INTRODUCTION

Quantum entanglement is the key resource for applications to quantum computation and quantum information theory. See Ref. 20. One of the most important topics in the theory of entanglement is how to distinguish entangled states from separable states. In the early eighties, Choi⁵ observed that if a positive (‘positive’ means ‘positive semi-definite’) matrix in the tensor product belongs to tensor product of positive parts then its partial transpose must be positive. Since the tensor product of positive parts is just the convex cone generated by separable states, this is equivalent to the PPT criterion which was rediscovered later by Peres²⁹. This criterion is already implicit in the Woronowicz’ earlier work⁴⁰ who initiated the duality theory between positivity of linear maps and the separability of states. He constructed an example of positive partial transposed (PPT) entangled state in order to show the existence of indecomposable positive linear maps.

It was Horodecki’s¹⁷ who used the relation between positive maps and separable states to get the complete criterion for separability. The Woronowicz’s idea has been developed for general cases in Ref. 10 to get the duality theory for s -positive linear maps. For the case of $s = 1$, this is equivalent to the Horodecki’s criterion, through the Jamiolkowski-Choi isomorphism^{4,21}. We note that the general cases give rise to the notion of Schmidt numbers and the relation to s -positivity of linear maps, which was obtained independently in Ref. 31 and 36. See also Refs. 32 and 33 for approaches to the duality theory which works for an arbitrary mapping cone. Horodecki’s criterion tells us that we need positive maps in order to detect entanglement, which has been reformulated in terms of entanglement witnesses³⁵. Under the JC isomorphism, an entanglement witness is just a positive linear map which is not completely positive. An entanglement witness which detects a maximal set of entanglement is said to be optimal, as was introduced in Ref. 26.

In spite of its importance, the whole structure of the convex cone of all positive linear maps is far from being completely understood, and there had even been very few known nontrivial examples of positive linear maps until the eighties. To overcome this difficulty, the idea to consider the line segment from the trace map to a given positive map had been used in mathematical literature to distinguish various notions of positivity. See Ref. 37 and 38. From the point of view of physics, positive maps which are not completely positive, like the transpose maps, are non-physical operations. The main idea of structural physi-

cal approximation (SPA)^{11,18,19} is to approximate positive maps by the nearest completely positive maps, which are physical operations, and so we can implement them in the real world. It was theoretically shown²² and later demonstrated in practice^{27,28} that the SPA's of the transpose map and the partial transpose map are experimentally feasible. On the mathematics side, SPAs correspond to the points at which a segment between the trace map and a given positive map crosses the border of the set of completely positive maps.

We recall that positive maps (respectively completely positive maps) correspond to block-positive (respectively positive) matrices in $M_m \otimes M_n$ under the JC isomorphism, where M_n denotes the $*$ -algebra of all $n \times n$ matrices over the complex field, with the identity $\mathbb{1}_n$. We also note that the trace map corresponds to $\mathbb{1}_m \otimes \mathbb{1}_n$. For an entanglement witness $W \in M_m \otimes M_n$, we consider the line segment L_W from $\mathbb{1}_m \otimes \mathbb{1}_n$ to W . The SPA of W is the positive matrix on L_W nearest to W . It was conjectured in Ref. 22 that if W is an optimal entanglement witness then its SPA is separable. Several authors^{1,6-9,30} considered various classes of optimal entanglement witnesses to support the conjecture. Here, we provide a counterexample. Since non-decomposable optimal witnesses need not be optimal in the sense of Section IV of Ref. 26, we use the term PPTES witnesses¹⁵. Here, PPTES refers to PPT entangled states. PPTES optimality of a witness W means that W detects a maximal set (in the sense of inclusion) of PPT entangled states. This is equivalent to say that W is indecomposable and both W and W^Γ is optimal²⁶, where W^Γ denotes the partial transpose of W . Let us recall that a special role is played by the product vectors $\phi \otimes \psi$ that satisfy $\langle \phi \otimes \psi | W | \phi \otimes \psi \rangle = 0$. If these vectors span $\mathbb{C}^m \otimes \mathbb{C}^n$, we say that W has the *spanning property*. We say that W is co-optimal (respectively co-spanning) if W^Γ is optimal (respectively spanning), bi-optimal if it is both optimal and co-optimal, and bi-spanning similarly.

In order to deal with the SPA conjecture in a systematic way, we introduce the notions of positive type and copositive type for entanglement witnesses in the next section, and show that if the SPA of an entanglement witness is separable, then the witness must be of copositive type. We also see that W is of positive type if and only if W^Γ is of copositive type. Because the optimality of PPTES witnesses is invariant under the operation of partial transpose, we conclude that only approximately half of optimal PPTES witnesses can satisfy the SPA conjecture. In the third section, we exhibit examples to show that the SPA conjecture fails even for the case of copositive type. We consider entanglement witnesses

given in Ref. 16 to find examples of optimal PPTES witnesses of copositive type whose SPA are not separable. The SPA's of our examples are exactly $3 \otimes 3$ PPT edge states²⁵ of type (6, 8). Very recently, after the authors posted this paper, Størmer³⁴ also examined the same class of entanglement witnesses¹⁶ to find optimal indecomposable entanglement witnesses whose SPA are not separable.

II. POSITIVE AND COPOSITIVE TYPE.

From now on, we say that a block matrix is *copositive* if its partial transpose is positive. Copositive matrices correspond to completely copositive maps under the JC isomorphism. For a given block-positive matrix W , we consider the line segment L_W from $\mathbb{1}_n \otimes \mathbb{1}_m$ to W , and compare the distances to the nearest positive matrix and the nearest copositive matrix on L_W . We say that W is of *positive* (resp. *copositive*) *type* if the distance to the positive (resp. copositive) part is shorter than or equal to the other. We also say that a block-positive matrix is of *PPT type* if two distances coincide.

To be precise, we consider self-adjoint matrix $W_t = (1 - t)/(mn)\mathbb{1}_m \otimes \mathbb{1}_n + tW$. When t_0 is the largest number in the interval $[0, 1]$ for which W_{t_0} is positive, W_{t_0} is said to be the SPA of W . It is clear that the SPA of W is separable if and only if the following condition

$$0 \leq t \leq 1, W_t \text{ is positive} \implies W_t \text{ is separable} \quad (1)$$

holds. It is also clear that the validity of the condition (1) is not changed when we replace $1/(mn)\mathbb{1}_m \otimes \mathbb{1}_n$ in the definition of W_t by $\mathbb{1}_m \otimes \mathbb{1}_n$ alone due to the convexity. Therefore, we may redefine W_t as

$$W_t := (1 - t)\mathbb{I}_m \otimes \mathbb{I}_n + tW$$

for every block-positive W , so far as we are concerned with the SPA conjecture.

Since separable states are of PPT, we see that if the SPA conjecture is true then every optimal entanglement witness W must satisfy the following condition

$$0 \leq t \leq 1, W_t \text{ is positive} \implies W_t \text{ is of PPT}$$

which is weaker than (1), and so satisfies the following equivalent condition

$$0 \leq t \leq 1, W_t \text{ is positive} \implies W_t \text{ is copositive.} \quad (2)$$

We define two real numbers α_W and β_W in the interval $[0, 1]$ by

$$\begin{aligned}\alpha_W &:= \sup\{t \in [0, 1] : W_t \text{ is positive}\}, \\ \beta_W &:= \sup\{t \in [0, 1] : W_t \text{ is copositive}\},\end{aligned}$$

for an arbitrary block-positive W . We see that both α_W and β_W are nonzero, since $\mathbb{I}_m \otimes \mathbb{I}_n$ is an interior point of the convex cone generated by all separable states. We also note that the number $1 - \alpha_W$ (resp. $1 - \beta_W$) plays a role of the distance from W to the positive (resp. copositive) part through the line segment L_W . It is clear that W satisfies the condition (2) if and only if the inequality $\alpha_W \leq \beta_W$ holds. Since W_t is positive if and only if W_t^Γ is copositive, we also have $\alpha_W = \beta_{W^\Gamma}$ and $\alpha_{W^\Gamma} = \beta_W$. Therefore, we see that $\alpha_W < \beta_W$ if and only if $\alpha_{W^\Gamma} > \beta_{W^\Gamma}$. In other words, W satisfies the condition (2) if and only if W^Γ does not satisfy (2), whenever $\alpha_W \neq \beta_W$.

We see that a block-positive matrix W is of positive type if and only if $\alpha_W \geq \beta_W$, and of copositive type if and only if $\alpha_W \leq \beta_W$ if and only if the condition (2) holds. We also see that W is of PPT type if and only if $\alpha_W = \beta_W$. The above discussion tells us that W is of positive type if and only if W^Γ is of copositive type, and the SPA conjecture can only hold for witnesses of copositive type. Especially, if the SPA of W is separable, then W is of copositive type and the SPA of W^Γ is never separable unless W is of PPT type.




FIG. 1. W has the mirror image W^Γ . W is nearer to the copositive part if and only if W^Γ is nearer to the positive part.

III. EXAMPLES OF PPT TYPE.

In this section, we exhibit examples of indecomposable entanglement witnesses of PPT type with the bi-spanning property in the sense of Ref. 15, whose SPAs are not separable.

For nonnegative real numbers a, b, c and $-\pi \leq \theta \leq \pi$, we consider the following self-adjoint block matrix in $M_3 \otimes M_3$:

$$W[a, b, c; \theta] = \begin{pmatrix} a & \cdot & \cdot & \cdot & -e^{i\theta} & \cdot & \cdot & \cdot & -e^{-i\theta} \\ \cdot & c & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & b & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & b & \cdot & \cdot & \cdot & \cdot & \cdot \\ -e^{-i\theta} & \cdot & \cdot & \cdot & a & \cdot & \cdot & \cdot & -e^{i\theta} \\ \cdot & \cdot & \cdot & \cdot & \cdot & c & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & c & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & b & \cdot \\ -e^{i\theta} & \cdot & \cdot & \cdot & -e^{-i\theta} & \cdot & \cdot & \cdot & a \end{pmatrix}.$$

We also put $p_\theta = \max\{q_{(\theta-\frac{2}{3}\pi)}, q_\theta, q_{(\theta+\frac{2}{3}\pi)}\}$, where $q_\theta = e^{i\theta} + e^{-i\theta}$. We note that $1 \leq p_\theta \leq 2$. We also see that $p_\theta = 2$ if and only if $\theta = 0, \pm 2\pi/3$, and $p_\theta = 1$ if and only if $\theta = \pm\pi/3, \pm\pi$. It is easy to see that $W[a, b, c; \theta]$ is of PPT if and only if

$$a \geq p_\theta, \quad bc \geq 1. \quad (3)$$

If $1 < p_\theta < 2$, the cases of $a = p_\theta$ and $bc = 1$ give rise to new kinds of examples²⁵ of PPT edge states of type $(8, 6)$. If $\theta = 0$ then the notion of PPT coincides²⁵ with separability. On the other hand, the authors¹⁴ explored the boundary structures between the separability and the inseparability for PPT states when $\theta = \pi$. We recall that the case $\theta = 0$ had been considered in Ref. 3.

It turns out¹⁶ that $W[a, b, c; \theta]$ is block-positive if and only if the condition

$$a + b + c \geq p_\theta, \quad a \leq 1 \implies bc \geq (1 - a)^2 \quad (4)$$

holds. We refer to Ref. 16 for the pictures of the 3-dimensional convex bodies determined by (3) and (4) for fixed θ . From now on, we concentrate on the case $a < p_\theta$ and $bc < 1$, for which $W[a, b, c; \theta]$ is neither positive nor copositive. We have

$$W_t[a, b, c; \theta] = t W \left[\frac{a_t}{t}, \frac{b_t}{t}, \frac{c_t}{t}; \theta \right], \quad 0 < t \leq 1,$$

with $a_t = 1 - t + ta$, $b_t = 1 - t + tb$, and $c_t = 1 - t + tc$. We see that $W_t[a, b, c; \theta]$ is positive if and only if $a_t \geq tp_\theta$ if and only if $t \leq \alpha_W = 1/(p_\theta + 1 - a)$, and $W_t[a, b, c; \theta]$ is copositive

if and only if $b_t c_t \geq t^2$ if and only if

$$F(t) := (b + c - bc)t^2 - (b + c - 2)t - 1 \leq 0.$$

Therefore, we see that $W[a, b, c; \theta]$ is of copositive type if and only if $0 \leq t \leq \alpha_W$ implies $F(t) \leq 0$. Since $F(0) = -1 < 0$ and $F(1) = 1 - bc > 0$, we see that this happens if and only if $F(\alpha_W) \leq 0$ if and only if

$$(p_\theta - a + b)(p_\theta - a + c) \geq 1.$$

Analogously, $W[a, b, c; \theta]$ is of positive type if and only if the reverse inequality holds, and of PPT type if and only if the equality holds.

On the other hand, we note that the SPA of $W[a, b, c; \theta]$ is given by

$$W_{\alpha_W}[a, b, c; \theta] = \frac{1}{p_\theta + 1 - a} W[p_\theta, p_\theta - a + b, p_\theta - a + c; \theta].$$

We note²⁵ that $W_{\alpha_W}[a, b, c; \theta]$ is not separable when

$$(p_\theta - a + b)(p_\theta - a + c) = 1, \quad 1 < p_\theta < 2, \quad (5)$$

that is when $W[a, b, c; \theta]$ is of PPT type. We concentrate on the following two subcases:

$$2 - p_\theta \leq a < 1, \quad a + b + c = p_\theta, \quad bc = (1 - a)^2, \quad (6)$$

$$1 \leq a < p_\theta, \quad a + b + c = p_\theta, \quad bc = 0. \quad (7)$$

We note¹⁶ that $W[a, b, c; \theta]$ is indecomposable in the above cases; has the bi-spanning property in the case (6) and has the co-spanning property in the case (7). See Ref. 12 and 13 for the case of $\theta = 0$. We also note that condition (5) together with (6) is translated into

$$3a^2 - 2(2p_\theta + 1)a + 2p_\theta^2 = 0, \quad 2 - p_\theta \leq a < 1,$$

and it is easy to see that

(i) There exists a, b and c satisfying (5) and (6) if and only if $4/3 \leq p_\theta < 1 + 1/\sqrt{2}$.

In this case, there are two solutions as is seen in FIG 2 when $2 - p_\theta < a < 1$. If $a = 2 - p_\theta$ then we have only one solution with $b = c = p_\theta - 1$. For the case (7), we note that the condition (5) is equivalent to

$$2(p_\theta - a)^2 = 1, \quad 1 \leq a < p_\theta < 2,$$

and thus we have the following:




FIG. 2. Figure of the boundary of the convex body determined by (3) and (4), which is lying on the plane $a + b + c = p_\theta$ for fixed θ . Continuous line in the case (i) (respectively (ii)) indicates a, b, c satisfying the condition (6) (respectively (7)). In both cases, black dots represent PPT types which violate the SPA conjecture. We refer to FIG. 3 of Ref. 16 for the pictures of the 3-dimensional convex bodies. The point A represents the case $a = 2 - p_\theta, b = c = p_\theta - 1$, and the point B represents the case $a = p_\theta, b = c = 0$. We refer to the figure in Ref. 23, Section 5 for the case of $\theta = 0$.

(ii) There exists a, b and c satisfying (5) and (7) if and only if $1 + 1/\sqrt{2} \leq p_\theta < 2$.

In the first case (i), we have indecomposable entanglement witnesses with the bi-spanning properties whose SPA are not separable. These entanglement witnesses are optimal PPTES witnesses in the sense of Ref. 15 which detect maximal sets of PPTES. In the case (ii), we see that the partial transposes W^Γ give rise to examples of optimal indecomposable entanglement witnesses whose SPA are not separable. It should be noted that they are not ‘nd-OWE’s in the sense of Ref. 26, or PPTES optimal witnesses using our terminology, when $1 + 1/\sqrt{2} < p_\theta < 2$, because they are not co-optimal; the smallest face determined by W^Γ contains $W[p_\theta, 0, 0; \theta]^\Gamma$ which is copositive. If $p_\theta = 1 + 1/\sqrt{2}$ then we have the solution $a = 1, b = p_\theta - 1, c = 0$ or $a = 1, b = 0, c = p_\theta - 1$, for which $W[a, b, c; \theta]$ are bi-optimal entanglement witnesses without the spanning property¹⁶.

IV. CONCLUSION.

In conclusion, we propose the notions of positive type and copositive type for block-positive matrices W . We have shown that if the SPA of W is separable then W must be of copositive type, and W is copositive type if and only if its partial transpose W^Γ is of positive type. Furthermore, a PPTES witness W is optimal if and only if W^Γ is optimal.

Therefore, using the above facts about positive and copositive types, we conclude that the separability of the SPA of an optimal PPTES witness W implies that the SPA of W^Γ is not separable, unless W is of PPT type. This is so, even though W^Γ is still an optimal PPTES witness. We provide concrete examples of indecomposable entanglement witnesses of both copositive type and PPT type which violate the SPA conjecture. It would be very nice to characterize entanglement witnesses whose SPA are separable.

Especially, it is still open if the SPA of an optimal decomposable entanglement witness is separable or not. Recent development^{2,24} on the optimality for decomposable cases might be helpful to determine if the SPA conjecture is true for decomposable case.

ACKNOWLEDGMENTS

KCH is partially supported by NRFK 2012-0002600. SHK is partially supported by NRFK 2012-0000939. This is a revised and expanded version of the preprint circulated under the same title, authored by SHK alone. Examples of PPT type are included in this revised version. The authors are grateful to Antonio Acin, Mafalda Almeida, Remigiusz Augusiak, Joonwoo Bae, Jaroslaw Korbicz and Maciej Lewenstein for their feedback on the preprint. Special thanks are due to Joonwoo Bae who informed us their discussion about the preprint, and Remigiusz Augusiak who pointed out errors in the preprint. The authors are also grateful to Hyang-Tag Lim for helpful discussion on the papers^{27,28}.

REFERENCES

- ¹R. Augusiak, J. Bae, Ł. Czekaj, and M. Lewenstein, “On structural physical approximations and entanglement breaking maps,” *J. Phys. A* **44**, 185308 (2011).
- ²R. Augusiak, G. Sarbicki, and M. Lewenstein, “Optimal decomposable witnesses without the spanning property,” *Phys. Rev. A* **84**, 052323 (2011).
- ³S.-J. Cho, S.-H. Kye, and S. G. Lee, “Generalized Choi maps in three-dimensional matrix algebra,” *Linear Algebra Its Appl.* **171**, 213–224 (1992).
- ⁴M.-D. Choi, “Completely positive linear maps on complex matrices,” *Linear Algebra Its Appl.* **10**, 285–290 (1975).

- ⁵M.-D. Choi, *Operator Algebras and Applications* (Kingston, 1980), pp. 583–590, Proc. Sympos. Pure Math. Vol 38. Part 2, Amer. Math. Soc., 1982.
- ⁶D. Chruściński, and J. Pytel, “Constructing optimal entanglement witnesses. II. Witnessing entanglement in 4N4N systems,” *Phys. Rev. A* **82**, 052310 (2010).
- ⁷D. Chruściński and J. Pytel, “Optimal entanglement witnesses from generalized reduction and Robertson maps,” *J. Phys. A* **44**, 165304 (2011).
- ⁸D. Chruściński, J. Pytel, and G. Sarbicki, “Constructing optimal entanglement witnesses,” *Phys. Rev. A* **80**, 062314 (2009), 062314.
- ⁹D. Chruściński and F. A. Wudarski, “Geometry of Entanglement Witnesses for Two Qutrits,” *Open Syst. Inf. Dyn.* **18**, 375–387 (2011).
- ¹⁰M.-H. Eom, and S.-H. Kye, “Duality for positive linear maps in matrix algebras,” *Math. Scand.* **86**, 130–142 (2000).
- ¹¹J. Fiurasek, “Structural physical approximations of unphysical maps and generalized quantum measurements,” *Phys. Rev. A* **66**, 052315 (2002).
- ¹²K.-C. Ha and S.-H. Kye, “One-parameter family of indecomposable optimal entanglement witnesses arising from generalized Choi maps,” *Phys. Rev. A* **84**, 024302 (2011).
- ¹³K.-C. Ha and S.-H. Kye, “Entanglement Witnesses Arising from Exposed Positive Linear Maps,” *Open Syst. Inf. Dyn.* **18**, 323–337 (2011).
- ¹⁴K.-C. Ha and S.-H. Kye, “Geometry of the faces for separable states arising from generalized Choi maps,” *Open Syst. Inf. Dyn.* **19**, 125009 (2012).
- ¹⁵K.-C. Ha and S.-H. Kye, “Optimality for indecomposable entanglement witnesses,” *Phys. Rev. A* (to appear), e-print arXiv:1204.6596.
- ¹⁶K.-C. Ha and S.-H. Kye, “Entanglement witnesses arising from Choi type positive linear maps,” e-print arXiv:1205.2921.
- ¹⁷M. Horodecki, P. Horodecki, and R. Horodecki, “Separability of mixed states: necessary and sufficient conditions,” *Phys. Lett. A* **223**, 1–8 (1996).
- ¹⁸P. Horodecki, “From limits of quantum operations to multicopy entanglement witnesses and state-spectrum estimation,” *Phys. Rev. A* **68**, 052101 (2003).
- ¹⁹P. Horodecki, and A. Ekert, “Method for Direct Detection of Quantum Entanglement,” *Phys. Rev. Lett.* **89**, 127902 (2002).
- ²⁰R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, “Quantum entanglement,” *Rev. Mod. Phys.* **81**, 865–942 (2009).

- ²¹A. Jamiolkowski, “An effective method of investigation of positive maps on the set of positive definite operators,” *Rep. Math. Phys.* **5**, 415–424 (1974).
- ²²J. K. Korbicz, M. L. Almeida, J. Bae, M. Lewenstein, and A. Acin, “Structural approximations to positive maps and entanglement-breaking channels,” *Phys. Rev. A* **78**, 062105 (2008), 062105.
- ²³S.-H. Kye, “Facial structures for various notions of positivity and applications to the theory of entanglement,” e-print arXiv:1202.4255.
- ²⁴S.-H. Kye, “Necessary conditions for optimality of decomposable entanglement witness,” *Rep. Math. Phys.*(to be published), e-print arXiv:1108.0456.
- ²⁵S.-H. Kye and H. Osaka, “Classification of bi-qutrit positive partial transpose entangled edge states by their ranks,” *J. Math. Phys.* **53**, 052201 (2012)
- ²⁶M. Lewenstein, B. Kraus, J. I. Cirac, and P. Horodecki, “Optimization of entanglement witnesses,” *Phys. Rev. A* **62**, 052310 (2000).
- ²⁷H.-T. Lim, Y.-S. Kim, Y.-S. Ra, J. Bae and Y.-H. Kim, “Experimental Realization of an Approximate Partial Transpose for Photonic Two-Qubit Systems,” *Phys. Rev. Lett.* **107**, 160401 (2011).
- ²⁸H.-T. Lim, Y.-S. Ra, Y.-S. Kim, J. Bae, and Y.-H. Kim, “Experimental implementation of the universal transpose operation using the structural physical approximation,” *Phys. Rev. A* **83**, 020301 (2011).
- ²⁹A. Peres, “Separability Criterion for Density Matrices,” *Phys. Rev. Lett.* **77**, 1413 (1996).
- ³⁰X. Qi, and J. Hou, “Characterization of optimal entanglement witnesses,” *Phys. Rev. A* **85**, 022334 (2012).
- ³¹A. Sanpera, D. Bruß, and M. Lewenstein, “Schmidt-number witnesses and bound entanglement,” *Phys. Rev. A* **63**, 050301 (2001).
- ³²L. Skowronek, “Cones with a mapping cone symmetry in the finite-dimensional case,” *Linear Algebra Its Appl.* **435**, 361–370 (2011).
- ³³E. Størmer, “Duality of cones of positive maps,” *Münster J. Math.* **2**, 299–310 (2009).
- ³⁴E. Størmer, “Separable states and SPA of a positive map,” e-print arXiv:1206.563.
- ³⁵B. M. Terhal, “Bell inequalities and the separability criterion,” *Phys. Lett. A* **271**, 319–326 (2000).
- ³⁶B. M. Terhal, and P. Horodecki, “Schmidt number for density matrices,” *Phys. Rev. A* **61**, 040301 (2000).

- ³⁷T. Takasaki and J. Tomiyama, “On the geometry of positive maps in matrix algebras,” Math. Z. **184**, 101–108 (1983).
- ³⁸J. Tomiyama, “On the geometry of positive maps in matrix algebras. II,” Linear Algebra Its Appl. **69**, 169–177 (1985).
- ³⁹J. Tomiyama, “On the geometry of positive maps in matrix algebras,” Contemp. Math. **62**, 357 (1987).
- ⁴⁰S. L. Woronowicz, “Positive maps of low dimensional matrix algebras,” Rep. Math. Phys. **10**, 165–183 (1976).